

Phenylephrine Hydrochloride Chlorpheniramine Maleate

Bronchofen™

Syrup / Drops

DECONGESTANT/ANTIHISTAMINE

FORMULATION:

SYRUP:

Each 5 mL (1 teaspoonful) contains:

Phenylephrine Hydrochloride	5 mg
Chlorpheniramine Maleate	1 mg

DROPS:

Each mL contains:

Phenylephrine Hydrochloride	1 mg
Chlorpheniramine Maleate	0.8 mg

PROPERTIES AND ACTIONS:

Bronchofen combines two widely used agents, phenylephrine HCl and chlorpheniramine maleate which provide relief of symptoms such as nasal congestion, watery eyes, sneezing, itchiness and runny nose that accompany common cold, sinusitis and allergic rhinitis.

Phenylephrine, a decongestant, relieves nasal congestion associated with common cold, hay fever/allergic rhinitis, sinusitis and other respiratory diseases.

Chlorpheniramine maleate, an antihistamine, relieves itchiness and other clinical expressions of allergy.

INDICATIONS:

As supportive agent in the following clinical disorders:

- Common cold
- Allergic rhinitis
- Sinusitis
- Allergic conjunctivitis associated with rhinitis

DOSAGE:

SYRUP:

2 - 6 yrs. old:	2.5 mL (1/2 teaspoon)
7 - 12 yrs. old:	5 mL (1 teaspoon)
12 yrs. & older:	7.5 mL (1 1/2 teaspoons)

To be taken 3-4 times a day or as prescribed by the physician.

DROPS:

1 - 6 months.:	0.5 mL - 0.75 mL
6 - 12 months.:	0.75 mL - 1.0 mL
1 - 2 yrs.:	1.0 mL - 1.25 mL

To be taken 3-4 times a day or as prescribed by the physician.

SIDE EFFECTS:

Chlorpheniramine maleate may cause dizziness and sedation. Phenylephrine may cause dizziness, tremors and an increase in blood pressure. Mild gastrointestinal and central nervous system disturbances may also occur.

PRECAUTIONS/CONTRAINDICATIONS:

As chlorpheniramine maleate may cause drowsiness and sedation, patients must avoid engaging in activities which require mental alertness.

Phenylephrine-Chlorpheniramine (Bronchofen) should not be taken by asthmatic patients or by patients with known hypersensitivity to any of the active ingredients. It should not be administered concurrently with monoamine oxidase inhibitors or within 14 days of stopping such treatment.

Phenylephrine-Chlorpheniramine (Bronchofen) is contraindicated in patients with history of hypersensitivity to the active ingredients, in coronary artery disease, hypertension, hyperthyroidism, hyperexcitability, acute asthma attacks, tumor of the adrenal gland (pheochromocytoma) narrow-angle glaucoma and severe liver function impairment. It should be used with great caution by individuals with history of asthma, high blood pressure, diabetes, thyroid disease, glaucoma and enlargement of the prostate gland.

Phenylephrine, should be given with great caution in patients with diabetes mellitus. Safety in pregnancy and lactation has not been established.

CAUTION:

Foods, Drugs, Devices and Cosmetics Act prohibits dispensing without prescription.

AVAILABILITY:

Syrup: Bottles of 60 mL DR-XY29289

Drops: Bottles of 15 mL DR-XY29196

STORE AT TEMPERATURES NOT EXCEEDING 30OC.

“For suspected adverse drug reaction, report to the FDA: www.fda.gov.ph”

SVMORE

Manufactured for:

S.V. MORE PHARMA CORPORATION

5/F S.V. More Group Corporate Center

16 Scout Tuazon cor. Roces Ave., Quezon City

Metro Manila, Philippines

by: Lloyd Laboratories, Inc.

No.10 Lloyd Ave., First Bulacan Industrial City,

City of Malolos, Bulacan